

RESEARCH ARTICLE

Prevalence of Bovine Mastitis Pathogens in Bulk Tank Milk in China

Yanliang Bi¹, Ya Jing Wang¹, Yun Qin², Roger Guix Vallverdú², Jaime Maldonado García², Wei Sun², Shengli Li^{1*}, Zhijun Cao^{1*}

1 State Key Laboratory of Animal Nutrition, Beijing Engineering Technology Research Center of Raw Milk Quality and Safty Control, College of Animal Science and Technology, China Agricultural University, Beijing, China, **2** Laboratorios Hipra S.A., Amer Giron, Spain

* lisheng0677@163.com(SLL); caozhijun@cau.edu.cn (ZJC)



OPEN ACCESS

Citation: Bi Y, Wang YJ, Qin Y, Guix Vallverdú R, Maldonado García J, Sun W, et al. (2016) Prevalence of Bovine Mastitis Pathogens in Bulk Tank Milk in China. PLoS ONE 11(5): e0155621. doi:10.1371/journal.pone.0155621

Editor: Glenn F. Browning, The University of Melbourne, AUSTRALIA

Received: September 23, 2015

Accepted: May 2, 2016

Published: May 17, 2016

Copyright: © 2016 Bi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: The funder, China Agricultural University (grant number: CARS-37), had a role in study design, data collection and analysis, decision to publish, and preparation of the manuscript. The funder, Laboratorios Hipra S.A., had a role in study design, sample collection, materials preparation, and proofreading of the manuscript for mistakes and grammatical errors. Received the funding: YLB.

Competing Interests: The authors have declared that no competing interests exist. Four employees

Abstract

The objectives of this study were to estimate the herd prevalence of major mastitis pathogens in bulk tank milk (BTM) in China dairy herds, to determine the relationship between the presence of mastitis pathogens and bulk tank milk somatic cell counts (BTSCC), and to investigate the impact of different dairy cattle farming modes and region on bacterial species. BTM samples collected from 894 dairy herds in China were examined for the presence of mastitis pathogens. The Flinders Technology Associates (FTA) cards were used for BTM sample collection, storage, and transportation and bacterial DNA amplification by real-time PCR. Among contagious pathogens, *Staphylococcus aureus*, *Streptococcus agalactiae*, and *Streptococcus dysgalactiae* were detected in 50.1, 92.2, and 72.3% of the 894 BTM samples, respectively. Among environmental pathogens, *E. coli*, *Streptococcus uberis*, *Enterococcus* spp., *Klebsiella* spp., *Serratia marcescens*, *Corynebacterium bovis*, and *Arcanobacterium pyogenes* were detected in 28.6, 8.9, 35.7, 20.0, 1.3, 17.0, and 67.2% of the BTM samples, respectively. Staphylococcal β -lactamase gene was detected in 61.7% of the BTM samples. The presence of *Staphylococcus aureus* and *Arcanobacterium pyogenes* were significantly associated with high BTSCC, respectively. Significant differences were found in presence of *Staphylococcus aureus*, *Streptococcus agalactiae*, and *Streptococcus dysgalactiae* in BTM sampled from the small household farms, dairy-farming communities, and large-scaled dairy farms. There were significant differences in the presence of *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, *Arcanobacterium pyogenes*, staphylococcal β -lactamase gene, *Staphylococcus* spp., *Klebsiella* spp., *Enterococcus* spp., and *Streptococcus uberis* in BTM among Inner Mongolia, Heilongjiang, and Hebei province. In conclusion, contagious mammary pathogens are predominated among pathogens in BTM samples in China.

Introduction

Bovine mastitis is the inflammation of the mammary gland which can be caused by several bacterial species or species groups. Previous studies showed that the overwhelming majority of clinical and subclinical mastitis were caused by approximately 10 bacterial species or species

from the commercial funder Laboratorios Hipra S.A., Yun Qin, Roger Guix Vallverdú, Jaime Maldonado García, and Wei Sun, were listed as authors in the article due to their participant work to the current research. Yun Qin and Wei Sun had a role in sample collection. Roger Guix Vallverdú had a role in study design, data analysis, and proofreading of the manuscript. Jaime Maldonado García had a role in data analysis and materials preparation. The authors declare that no competing interests exist relating to employment, consultancy, patents, products in development, or marketed products. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials.

groups [1–3]. Mastitis pathogens are broadly classified as contagious and environmental pathogens. *Staphylococcus aureus*, *Streptococcus agalactiae*, and *Streptococcus dysgalactiae* are considered as contagious pathogens, which adapt to the environment of mammary gland and can potentially be spread from cow to cow during milking. *Streptococcus uberis*, *Enterococcus* spp., coliforms, *Arcanobacterium pyogenes*, and coagulase-negative staphylococci (CNS) are considered as environmental pathogens, which are described as opportunistic pathogens of the mammary gland and can potentially be transferred from the contaminated environment to the mammary gland of a cow during milking [1, 4].

Early identification of the prevalence and distribution of causative pathogens is of the most importance to effectively prevent diseases and to guide treatment [5–7]. Bulk tank milk (BTM) analysis has been used as a useful monitoring tool to assess milk quality and mastitis pathogens for many years [8–10]. The BTM sample test is less expensive, faster, and more convenient compared with testing quarter milk samples [11–12]. However, in contrast to quarter milk sample test, results of BTM sample test not only reveal the presence of mammary pathogens in a herd but also reflect milk contamination from the teat surface and the environment [1, 7].

The real-time PCR, which has predefined cycle threshold (Ct) values and internal controls allowing differentiation between truly positive and false negative results, has proved useful to identify mastitis pathogens [13–14]. It is based on identification of bacterial genome sequences, and can detect growth-inhibited and dead bacteria within 3 to 4 hours [14]. The real-time PCR assay has high sensitivity and specificity for diagnosis of mastitis pathogens in BTM. Mweu et al. [15] reported that, at a cycle threshold cut-off value of 40, the sensitivity and specificity of the real-time PCR used for screenings of BTM samples for *S. agalactiae* were 95.2 and 98.8%, respectively.

Flinders Technology Associates (FTA) cards technique is a well-established method for collection and storage of bacterial DNA samples [16–17]. FTA cards comprise cotton-based cellulose paper, impregnated with proprietary chemicals that serve to lyse cells, denature proteins, stabilize and preserve nucleic acids within the matrix fibre. By doing so the nucleic acids in the samples are immobilized thus allowing to transport easily and long-term storage at room temperature [18]. FTA cards have been used for preparation of DNA or RNA from a variety of biological materials, including blood, saliva, animal and plant tissues, and various pathogens coming from human, animal and plant samples [19–23]. In a previous report [24], raw and mastitis milk, as well as dairy process and product samples were tested using both FTA cards and PCR to detect *S. aureus* among other bacteria.

A considerable number of studies of mastitis pathogens in BTM have been published internationally [1, 3, 10]. To our knowledge, there are no scientific reports on the prevalence of major mastitis pathogens in BTM in China. The objectives of this study were 1) to use real time PCR to estimate the prevalence of mastitis pathogens in BTM from Chinese dairy herds, 2) to determine the relationships between mastitis pathogens and bulk tank milk somatic cell counts (BTSCC), and 3) to investigate the effect of geographic region and different dairy cattle farming modes on bacterial species prevalence in BTM.

Materials and Methods

Participants

All experimental protocols were reviewed and approved by the Animal Care Committee of China Agricultural University. Experimental animals care and use for this experiments were performed following the established guidelines of the College of Animal Science and Technology, China Agricultural University.

The study was conducted in the top 3 milk-producing provinces (Hebei, Heilongjian and Inner Mongolia) of China. In 2012, 52.3% of the total number of dairy cattle and 45.6% of the

total milk output in China came from these 3 provinces (National Bureau of Statistics of China 2013). The 3 provinces are all located in the northern region of China and also have the longest history of dairy farming among all the Chinese provinces. Hebei province (E 113°27'~119°50', N 36°05'~42°40') is located in the eastern plain of China. Heilongjiang province (E 121°11'~135°05', N 43°25'~53°33') belongs to the plain of northern cold region. Inner Mongolia (E 97°10'~126°02', N 37°30'~53°20') is located in the plateau region of northwest China. Herd selection was supported by provincial bureau of animal husbandry of the three provinces who provided the dairy herds information. 1,000 herds were randomly selected from the three provinces (400 herds from Mongolia, 300 herds from Heilongjiang province, 300 herds from Hebei province) and were invited to participate in the study. Dairy producers who participated in the study were solicited by provincial bureau of animal husbandry through their announcements about the study. Dairy producers who provided written response to the invitation were included in the study. A total of 894 dairy herds from Inner Mongolia (375 herds), Heilongjiang (243 herds) and Hebei (275 herds) responded in writing to the invitation and participated in this study (with approximately 228,000 lactating cows). The proportion of lactating cows in this study are 6.5% in Inner Mongolia, 5.9% in Heilongjiang, and 8.4% in Hebei, respectively. The average lactating cow herd size was 255 cows and ranged from 5 to 6563 cows. In China, BTSCC and bacteria counts are routinely analyzed by milk processing factories which collect milk from dairy farms/communities. Out of the 894 dairy herds, 311 herds (161 herds from Mongolia, 91 herds from Heilongjiang province, 59 herds from Hebei province) were not willing to share their data of BTSCC. Therefore, a total of 583 dairy herds' data of BTSCC were collected.

The dairy cattle farming modes were classified into three types according to their scale and organization in China, including small household farms, dairy-farming communities (cooperative dairy farms) and large-scaled dairy farms. Small household farm refers to the family farm which normally has less than 30 dairy cattle, which are not grouped, kept in the backyard and milked at the milk collection station close by. This dairy cattle farming mode is relatively simple and the milk yield on the farm is low. The dairy-farming community is a transitional form between the small household farm and the large-scaled dairy farm. In general, the number of dairy cattle is greater than 100 in a dairy-farming community. The dairy cattle owned by different farmers are raised in different barns on one community farm and separated according to ownership rather than grouped and raised according to the age and the physiological cycle. In a dairy-farming community, different dairy cattle belonged to different farmers are milked in one milking parlour and unified management methods can not be achieved. A large-scaled dairy farm refers to a dairy farm owning over 200 dairy cattle, where the cattle are grouped, bred and managed according to their ages and physiological cycles. Unified management methods are applied in this dairy cattle farming mode. Relatively, the large-scaled dairy farms are better mechanized, and raw milk quality and average milk yield are higher than that of other two dairy cattle farming modes. In this study, each milk collection station served as one dairy herd and provides one BTM sample, so did each dairy-farming community. Among the 894 dairy herds, the proportion of BTM samples came from small household farms, dairy-farming communities and large-scaled dairy farms were 47.9, 25.3, and 26.8%, respectively.

BTM sample and data collections

In the period of April 2012 to October 2013, a total of 894 BTM samples from Mongolia, Heilongjiang and Hebei provinces were collected. Sampling was conducted by 3 trained university students who followed a standardized protocol, and used a standard collection kit provided in advanced. Briefly, the milk in the bulk tank was mixed thoroughly before collection, and a

sample was taken from the top of the bulk tank using gloves and a sterile disposable plastic pipet. Approximately 100 μ l of each sample were spotted onto the matrix area of a FTA card aseptically and were allowed to air-dry for 1 h at room temperature. After air-dried, FTA cards were placed in hermetic polythene bags containing a desiccant, and were submitted to the molecular biology laboratory (Laboratorios Hipra S.A., Beijing, China).

The information of sampling sites, dairy cattle farming modes and BTSCC of herds were recorded. In China, BTSCC is routinely analyzed by milk processing factories. In this study, BTSCC was determined with a Fossomatic cell counter instrument (Foss Electric A/S, Hillerød, Denmark).

Milk sample preparation and DNA extraction

For DNA elution, the entire inoculation area of each FTA card (round shape) was removed and cut into small fragments using sterile scissors and forceps. To avoid cross-contamination, the scissors and forceps used to handle the inoculation were changed between samples and were disinfected with high temperature alcohol burner. The small fragments were transferred in a 2 ml sterile polypropylene tube with safety latch. 1 ml of PCR water was transferred to each tube containing samples, capped and vortexed for 15 seconds, and then incubated for 20 minutes in a dry block at 100°C. After incubation, the tube was vortexed for 15 seconds, centrifuged at low speed, and thus the supernatant was obtained.

Liquid samples (supernatants) were processed as if they had been fresh milk samples. DNA in each sample was extracted and purified using the spin column-based nucleic acid purification reagents and the manufacturer's instructions included in the assay kit (PathoProof™ Mastitis Complete-12 KingFisher™ Kit, Thermo Fisher Scientific, Beijing, China). The extracted DNA was stored at -20°C until further analysis.

Real-Time PCR analysis

The Real-Time PCR tests were performed using the reagents and protocol instructions included in the PCR assay kit as described by Koskinen et al. [14]. The PCR assay can simultaneously identify the following 12 bacterial species or species groups: *Staphylococcus aureus*, *Staphylococcus* spp. (including *Staphylococcus aureus* and coagulase-negative staphylococci), staphylococcal β -lactamase gene, *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, *Streptococcus uberis*, *Escherichia coli*, *Enterococcus* spp. (including *E. faecalis* and *E. faecium*), *Klebsiella* spp. (including *K. oxytoca* and *K. pneumoniae*), *Corynebacterium bovis*, *Serratia marcescens*, *Arcanobacterium pyogenes* and/or *Peptoniphilus indolicus*. The assay included 4 separate multiplex real-time PCR reactions, each detecting 3 bacterial species or species groups. Combination of the 4 different real-time PCR reactions results was the final result of one sample. The PCR test for the species-specific detection of the mentioned microorganisms has been shown to be suitable and to perform well, even when phylogenetically closely related non-target species were tested [14]. The assay included enough controls for the validation of each run including negative DNA extraction controls, internal amplification standard (positive PCR controls), and non-template control that were included in each of the 4 real-time PCR reactions.

The PCR mixture consisted of 10 μ l of the Master Mix, 5 μ l of Primer Mix 1/2/3/4 (including species-specific primers and probes), and 10 μ l of extracted DNA. The real-time PCR instrument thermal cycler Stratagene Mx3005P (Agilent Technologies Inc., Santa Clara, CA) was used for amplification. The amplification conditions were as follow: 95°C for 10 min, 1 cycle; 95°C for 5 s and 60°C for 1 min, 40 cycles.

The real-time PCR fluorescence threshold was set, according to the assay's Manual. When a Ct value was below 37, the reaction was positive. On the contrary, the reaction was negative. Positive and negative controls, and negative PCR control were included in each multiplex PCR reaction, for validation of the extraction and amplification process.

Statistical analysis

The date of the occurrence of 11 mastitis pathogens and the staphylococcal β -lactamase gene in BTM samples and BTSCC were transferred to Microsoft Excel. We defined pathogen-positive by real-time PCR assay as "1" and pathogen-free by real-time PCR assay as "0". All statistical analyses were performed using SAS 9.3 (SAS Inst. Inc., Cary, NC). Data were analyzed as a factorial arrangement using PROC GLMMIX procedure of SAS with fixed effects of province and dairy cattle farming mode. The correlation between the BTSCC and the presence of mastitis pathogens were analyzed using PROC CORR procedure of SAS. All results were reported as least-square means (LSM). BTM sample was the experimental unit. The statistical model used as follow,

$$Y_{ijk} = \mu + \gamma_i + \delta_j + (\gamma\delta)_{ij} + \epsilon_{ijk},$$

where Y_{ijk} was the proportion of the existence of mastitis pathogen, μ was overall mean, γ_i was province effect, δ_j was the dairy cattle farming mode effect, $(\gamma\delta)_{ij}$ was the interaction between province effect and dairy cattle farming mode effect, and ϵ_{ijk} was random error. A P -value less than 0.0042 (0.05/12) was considered a significant correlation between the BTSCC and the presence of mastitis pathogens. Considering the repeated analysis, differences among effects were declared significant at $P < 0.0042$.

Results

BTM samples from 894 dairy herds in 3 Chinese provinces were examined for mastitis pathogens. Proportion of BTM samples that were positive for 12 mastitis pathogen species or species groups are presented in Table 1. Among contagious pathogens, *S. aureus*, *S. agalactiae*, and *S. dysgalactiae* were detected in 448 (50.1%), 824 (92.2%), and 647 (72.3%) samples, respectively. Among environmental pathogens, *E. coli*, *S. uberis*, *Enterococcus* spp., *Klebsiella* spp., *S.*

Table 1. Proportion of 894 bulk milk samples that were positive for bovine mastitis pathogens in China from April 2012 to October 2013.

Bacterial species or species groups	Number of positive samples (herds)	Positive proportion of samples (%)
<i>S. aureus</i>	448	50.1
<i>S. agalactiae</i>	824	92.2
<i>S. dysgalactiae</i>	647	72.3
<i>E. coli</i>	255	28.6
<i>S. uberis</i>	79	8.9
<i>Enterococcus</i> spp.	319	35.7
<i>Klebsiella</i> spp.	178	20.0
<i>Serratia marcescens</i>	12	1.3
<i>C. bovis</i>	152	17.0
<i>A. pyogenes</i>	601	67.2
<i>Staphylococcus</i> spp.	850	95.1
staphylococcal β -lactamase gene	551	61.7

doi:10.1371/journal.pone.0155621.t001

marcescens, *C. bovis*, and *A. pyogenes* were detected in 255 (28.6%), 79 (8.9%), 319 (35.7%), 178 (20.0%), 12 (1.3%), 152 (17.0%), and 601 (67.2%) samples, respectively. 850 (95.1%) samples were tested positive for *Staphylococcus* spp., 551 (61.7%) samples were tested positive for staphylococcal β -lactamase gene, and 343 (38.4%) BTM samples were simultaneously detected positive for *S. aureus* and staphylococcal β -lactamase gene. Overall, 743 BTM samples (83.1%) out of the total 894 BTM samples were positive for 4 or more species of mastitis bacterial pathogens.

The geometric mean BTSCC of the 583 herds was 650,000 cells/ml. *S. agalactiae* was detected in almost all of the 583 BTM samples (97.9%). A significant correlation ($P < 0.0001$) was observed between the presence of *S. aureus* and high BTSCC (Table 2). There was a significant difference ($P = 0.0019$) in BTSCC between *S. aureus*-positive herds and -negative herds. The geometric mean BTSCC of *S. aureus*-positive herds was 595,000 cells/ml, which was higher than that of -negative herds (441,000 cells/ml). It was worth noting that a significant relationship ($P = 0.0039$) was found between *A. pyogenes*-positive herds and high BTSCC. A significant correlation ($P = 0.0007$) was found between the presence of staphylococcal β -lactamase gene and high BTSCC. There was no correlation between the presence of other pathogens and BTSCC ($P > 0.0042$). No difference was observed in BTSCC between other pathogens positive and negative herds ($P > 0.0042$).

There were significant differences ($P < 0.0001$) in the presence of *S. aureus*, *S. agalactiae*, and *S. dysgalactiae* in BTM between samples collected from the small household farms, dairy-farming communities, and large-scaled dairy farms (Table 3). The prevalence levels of the three mastitis pathogens in dairy-farming communities were higher than those on small household farms and large-scaled dairy farms. The dairy herd prevalence of mastitis-causing bacteria

Table 2. Correlations between the presence of mastitis pathogens and the mean bulk tank milk somatic cell counts, and differences in bulk tank milk somatic cell counts between the mastitis pathogens positive and negative herds.

Bacterial species or species groups	Number of herds (583)		Mean BTSCC ^a ($\times 10^4$ /ml)		Correlation Coefficients	P1 ^d	P2 ^e
	Positive ^b	Negative ^c	Positive ^b	Negative ^c			
<i>S. aureus</i>	310	273	59.5	44.1	0.33643	< .0001*	0.0019*
<i>E. coli</i>	111	472	54.2	49.5	0.03696	0.6384	0.366
<i>Staphylococcus</i> spp.	529	54	49.4	54.3	0.10149	0.196	0.5561
<i>Klebsiella</i> spp.	57	526	55.7	47.9	0.05035	0.522	0.2583
<i>Serratia marcescens</i>	9	574	40.6	63.1	-0.12319	0.1161	0.1659
<i>S. agalactiae</i>	574	9	59.1	44.5	0.15046	0.0545	0.4636
<i>Enterococcus</i> spp.	142	441	49.6	54.1	-0.06643	0.398	0.3421
<i>S. uberis</i>	77	506	53.6	50.0	0.12429	0.1128	0.5521
<i>S. dysgalactiae</i>	543	40	46.2	57.5	0.05295	0.5007	0.223
<i>C. bovis</i>	128	455	51.6	52.1	0.06596	0.4014	0.9196
<i>A. pyogenes</i>	486	97	59.7	44.0	0.22438	0.0039*	0.0185
β -lactamase ^f	358	225	54.9	48.8	0.26074	0.0007*	0.2226

^aBTSCC = bulk tank milk somatic cell counts.

^bPositive: positive by real-time PCR assay.

^cNegative: negative by real-time PCR assay.

^dP1: correlations *P* value.

^eP2: differences *P* value.

^f β -lactamase = staphylococcal β -lactamase gene.

* $P < 0.0042$.

doi:10.1371/journal.pone.0155621.t002

Table 3. Effect of dairy cattle farming modes on the presence of mastitis pathogens.

Bacterial species or species groups	SHF ^a	DFC ^b	LSDF ^c	SEM ^d	P
<i>S. aureus</i>	0.5199 ^f	0.6593	0.3804	0.0469	< .0001*
<i>E. coli</i>	0.3232	0.2363	0.2718	0.0433	0.4314
<i>Staphylococcus</i> spp.	0.9802	0.9146	0.9157	0.0229	0.0890
<i>Klebsiella</i> spp.	0.3066	0.1895	0.2240	0.0380	0.1227
<i>Serratia marcescens</i>	0.0170	0.0044	0.0152	0.0117	0.7394
<i>S. agalactiae</i>	0.6101	0.9683	0.8126	0.0244	< .0001*
<i>Enterococcus</i> spp.	0.3733	0.2852	0.3284	0.0438	0.4330
<i>S. uberis</i>	0.1446	0.0886	0.1177	0.0295	0.4744
<i>S. dysgalactiae</i>	0.5797	0.8072	0.7589	0.0344	< .0001*
<i>C. bovis</i>	0.1634	0.1594	0.1397	0.0363	0.8652
<i>A. pyogenes</i>	0.7162	0.7585	0.6973	0.0413	0.5457
β -lactamase ^e	0.6776	0.6385	0.5259	0.0453	0.0298

^aSHF = small household farm.

^bDFC = dairy-farming community.

^cLSDF = large-scaled dairy farm.

^dSEM = standard error of the mean.

^e β -lactamase = staphylococcal β -lactamase gene.

^f0.5199 = the values in each column represent the proportion of the existence of mastitis pathogens

*P < 0.0042.

doi:10.1371/journal.pone.0155621.t003

showed significant geographic variation (Table 4). There were significant differences (P < 0.0042) in the presence of *S. agalactiae*, *S. dysgalactiae*, *A. pyogenes*, staphylococcal β -lactamase gene, *Staphylococcus* spp., *Klebsiella* spp., *Enterococcus* spp., and *S. uberis* in BTM samples among the 3 Chinese provinces.

Table 4. Effect of regions on the presence of mastitis pathogens.

Bacterial species or species groups	Hebei	Heilongjiang	Inner Mongolia	SEM ^a	P
<i>S. aureus</i>	0.4431 ^c	0.5492	0.5674	0.0366	0.1466
<i>E. coli</i>	0.2629	0.227	0.3414	0.0347	0.1415
<i>Staphylococcus</i> spp.	0.9413	0.8746	0.9945	0.0441	0.0008*
<i>Klebsiella</i> spp.	0.2129	0.1587	0.3486	0.0383	0.0009*
<i>Serratia marcescens</i>	0.0100	0.0238	0.0027	0.0416	0.4261
<i>S. agalactiae</i>	0.6243	0.7921	0.9746	0.0246	< .0001*
<i>Enterococcus</i> spp.	0.3216	0.2302	0.4351	0.0473	0.0031*
<i>S. uberis</i>	0.1702	0.146	0.03466	0.0118	0.0018*
<i>S. dysgalactiae</i>	0.7408	0.8762	0.5289	0.0298	< .0001*
<i>C. bovis</i>	0.1404	0.1635	0.1586	0.0437	0.9035
<i>A. pyogenes</i>	0.8956	0.7079	0.5685	0.0457	< .0001*
β -lactamase ^b	0.6375	0.4079	0.7966	0.0231	< .0001*

^aSEM = standard error of the mean.

^b β -lactamase = staphylococcal β -lactamase gene.

^c0.4431 = the values in each column represent the proportion of the existence of mastitis pathogens

*P < 0.0042.

doi:10.1371/journal.pone.0155621.t004

Discussion

In the current study, the multiplex real-time PCR assay in conjunction with FTA technique was used for the first time in China to estimate the herd prevalence of major mastitis pathogens in BTM from 894 China dairy herds. The relationships between the presence of mastitis pathogens and BTSCC, as well as the impacts of dairy cattle farming modes and region on bacterial species in BTM were investigated.

FTA Card Technique and Real-Time PCR Assay

Because the areas of origin of the milk samples analyzed in this study were a great distance from the lab where they were analyzed, and due to the difficulty of guaranteeing the cold chain during transport, samples were inoculated and transported in FTA cards. BTM samples collected using FTA cards are easy to transport and face fewer shipping restrictions between the collection site and the laboratory without the need for a cold chain or liquid media handling [18]. The storage of BTM samples on FTA cards are also at room temperature and no refrigerators or freezers are required, which significantly reduces storage costs and space. The manufacturer states that genomic DNA entrapped in FTA card matrix fibre which stored at room temperature is stable for 17 years [18]. The FTA cards are impregnated with special chemicals that inactivate infectious bacteria and viruses, reduce the potential biohazards of samples, and avoid cross-contamination between samples [16, 25]. As the FTA card contains proprietary chemicals that lyse cells and inactivate bacteria while preserving nucleic acids, BTM samples collected with FTA cards cannot be used for the culture and isolation of mastitis pathogens, and are suitable for molecular diagnostics. Furthermore, the FTA card is small (7.3 cm×3.8cm) and can be shipped in a standard letter mail envelope, which is difficult for liquid based medium. The FTA card technique simplifies the DNA extraction and purification procedures and is a promising tool for BTM samples collection, transportation, storage, and DNA preparation for mastitis pathogens.

The real-time PCR has proved to be a valuable diagnostic technique for improving bacterial identification and furthered benefits to shorten throughput time, to allow fully objective interpretation of data, and to quantify the number of microorganism in milk samples [13–14]. Previous studies revealed that the real-time PCR assay has higher sensitivity and specificity for diagnosis of mastitis pathogens compared with bacterial culture [15, 26–27]. As with all methodologies, the real-time PCR assay need thorough validation, particularly to avoid false positive or negative results due to the capability of detecting low copy numbers of bacteria originated from environment or inhibition of the reaction induced by components present in the sample. In the study of Koskinen et al., false positive results was detected in real-time PCR in the quarter milk samples originating from clinically healthy cows with a low milk SCC [28]. In addition, the real-time PCR assay is limited to detecting only the bacterial species for which a test kit has been designed and the clinical relevance of detecting bacterial nucleic acids instead of viable cells should be evaluated [28]. Despite all this, the real-time PCR assay is still a promising molecular method for detecting of mastitis pathogens.

This study used the PathoProof Mastitis Complete-12 KingFisher Kit (Finnzymes Oy, Espoo, Finland) to identify all major pathogens responsible for bovine mastitis from BTM samples. The PathoProof Mastitis PCR Assay, including all reagents for DNA extraction and the real-time PCR, is a commercially available PCR-based mastitis test. The assay can be used to extract DNA directly from raw or preserved milk, and it was developed to identify a total of 11 major mastitis-causing pathogens and the β -lactamase gene responsible for staphylococcal penicillin resistance [3, 14]. The assay was validated by Koskinen et al. [14], who demonstrated that the assay has 100% specificity and 100% sensitivity in identification of a large collection of culture strains originating from bovine mastitis. Taponen et al. [26] showed that the assay was

able to identify mastitis bacteria in clinical mastitis samples with no bacterial growth in conventional culture. The assay has been used for identification of mastitis pathogens in all Danish dairy herds in November 2009 to January 2010 [3]. One disadvantage of the kit is that the real-time PCR assay is limited to detecting only the bacterial species without detecting *Mycoplasma bovis*. The PathoProof Mastitis Major-3 Kit used for identifying three contagious major mastitis pathogens including *M. bovis*, *S. aureus* and *S. agalactiae*, was recently reported [29]. In a word, the real-time PCR assay used in this study proved to be a useful tool for accurate diagnosis of mastitis-causing pathogens in BTM at routine milk recordings.

Prevalence of Pathogens

This study indicated that contagious mammary pathogens are predominated among pathogens in BTM samples in China. Our findings for the dominance of contagious pathogens are not in accordance with following researches at herd level. Katholm et al. [3] tested BTM from all 4,258 Danish dairy herds using real-time PCR analysis. *Staphylococcus* spp. were found in all BTM, *S. uberis* in 95%, *C. bovis* in 90%, *Enterococcus* spp. in 78%, *A. pyogenes* and/or *P. indolicus* in 63%, and *E. coli* in 61%, whereas *S. agalactiae*, *S. aureus*, and *S. dysgalactiae* were found in 7, 91, and 86% of the herds. Rysanek et al. [1] cultured BTM Samples from 268 randomly selected herds and found dominance of environmental pathogens over the contagious pathogens in the Czech Republic. The isolation rates of *S. uberis*, *Enterococcus* spp., *E. coli*, *S. aureus*, *S. agalactiae*, and *S. dysgalactiae* in BTM samples were 18.3, 24.3, 19.0, 19.8, 5.6, and 17.9%, respectively. Phuektes et al. [6] analysed BTM from 42 herds in Australia with a multiplex PCR and reported that *S. uberis* was the most frequently present species in BTM, being found in 44% of samples, and *S. aureus*, *S. agalactiae*, and *S. dysgalactiae* were present in 16, 22 and 28% of BTM samples, respectively.

Association between BTSCC and the presence of mastitis pathogens

The geometric mean of the BTSCC of China milk are higher compared with other industrialized countries [1, 6, 10]. In this study, there was a significant correlation between the presence of *S. aureus* and the high mean BTSCC in BTM. This is in agreement with previous studies [1, 5, 10, 30–33]. Riekerink et al. [10] reported a significant association between the mean BTSCC and the isolation of *S. aureus* from BTM samples. Wilson et al. [33] found that separate milking for cows infected with *S. aureus* reduced BTSCC from 600,000/ml to 345,000/ml. However, other studies showed that the mean BTSCC and detection of *S. aureus* were not significantly associated [31, 34]. Koop et al. [34] thought that the little effect on BTSCC was caused by the relatively low prevalence of infection with *S. aureus*, although udder infections with *S. aureus* can strongly increase quarter milk SCC.

Several studies showed that there was no significant relationship between the presence of environmental pathogens and BTSCC [1, 32]. This is in accordance with our results. In this study, no significant relationship were observed between BTSCC and the environmental mastitis-causing pathogen except *A. pyogenes*. It's worth noting that a significant relationship was found between the *A. pyogenes*-positive samples and the high BTSCC. Our results are in accordance with the findings of Klossowska et al. [35] and Malinowski et al. [36], who reported that the high SCC was significantly associated with intramammary infections by *A. pyogenes*.

Differences in the prevalence of mastitis pathogens between dairy cattle farming modes

The dairy herd's prevalence of mastitis-causing bacteria differed between the small household farms, dairy-farming communities and large-scaled dairy farms in our study. Significant

differences ($P < 0.0001$) were found in the prevalence of *S. aureus*, *S. agalactiae*, and *S. dysgalactiae* in BTM between the different dairy cattle farming modes. The prevalence levels of the 3 mastitis pathogens in dairy-farming communities were higher than those on small household farms and large-scaled dairy farms. *S. agalactiae* and *S. dysgalactiae* occurred less frequently in small household farms than that in dairy-farming communities and large-scaled dairy farms, whereas *S. aureus* occurred less frequently in large-scaled dairy farms. Herd size and farm management practices had considerable influence on the prevalence of mastitis pathogens in BTM [5, 7, 37]. In dairy-farming communities, the dairy cattle were milked by owners of cows, therefore the healthy cows and infected cows were milked at the same time. This may transmit the contagious pathogens from infected cows to healthy cows via the milking machinery. Another possible reason for dairy-farming communities having higher prevalence levels of contagious pathogens may be due to higher levels of cattle movement. Herds that introduce new cattle could significantly increase the risk of being infected with contagious pathogens [37–38].

Differences in the Prevalence of Mastitis Pathogens between Provinces

The dairy herd's prevalence of mastitis-causing bacteria differed between provinces in our study. There were highly significant differences ($P < 0.0042$) in the prevalence of *S. agalactiae*, *S. dysgalactiae*, *A. pyogenes*, staphylococcal β -lactamase gene, *Staphylococcus* spp., *Klebsiella* spp., *Enterococcus* spp. and *S. uberis* in BTM between the 3 provinces. The differences are mainly caused by geographic variation. Riekerink et al. [39] found that the distribution of mastitis-causing bacteria showed substantial geographic variation on Canadian dairy farms. Although highly significant differences in the prevalence of mastitis-causing bacteria were observed between the 3 Chinese provinces, the most frequently detected mastitis-causing bacteria species were almost the same. In Inner Mongolia, *Staphylococcus* spp., *S. agalactiae*, *A. pyogenes*, and *S. aureus* were the most frequently detected bacteria from dairy herds, whereas in Hebei and Heilongjiang, the most often discovered pathogens were *Staphylococcus* spp., *S. dysgalactiae*, *S. agalactiae*, and *A. pyogenes*.

In this study, sampling BTM was performed only once. This may underestimated the prevalence of mastitis pathogens due to intermittent shedding. Multiple sampling of BTM from the same herd over a period of time is needed to determine the true prevalence of contagious mastitis pathogens [10, 40–41].

Conclusions

The current study revealed that contagious mammary pathogens predominated in BTM samples in China. *Staphylococcus* spp., *S. agalactiae*, *S. dysgalactiae*, *A. pyogenes*, staphylococcal β -lactamase gene, and *S. aureus* were the most frequently detected pathogens in BTM samples. The presence of *S. aureus* and *A. pyogenes* were significantly associated ($P < 0.0042$) with the high BTSCC. There was a significant difference ($P < 0.0042$) in BTSCC between *S. aureus*-positive herds and *S. aureus*-negative herds. Significant differences ($P < 0.0001$) were found in the presence of *S. aureus*, *S. agalactiae*, and *S. dysgalactiae* in BTM between the small household farms, dairy-farming communities, and large-scaled dairy farms. The prevalence levels of the 3 mastitis pathogens in dairy-farming communities were higher than those on small household farms and large-scaled dairy farms. There were significant differences ($P < 0.0042$) in the presence of *S. agalactiae*, *S. dysgalactiae*, *A. pyogenes*, staphylococcal β -lactamase gene, *Staphylococcus* spp., *Klebsiella* spp., *Enterococcus* spp., and *S. uberis* in BTM between Inner Mongolia, Heilongjiang, and Hebei Province.

Acknowledgments

The authors would like to acknowledge all participating dairy farms, milk processors, and the support of milk factories. Teachers, colleagues, and the staff of the Laboratorios Hipra S.A. who participated in the study are also acknowledged.

Author Contributions

Conceived and designed the experiments: SLL ZJC RGV. Performed the experiments: YLB WS. Analyzed the data: YLB RGV JMG. Contributed reagents/materials/analysis tools: YQ JMG WS. Wrote the paper: YLB RGV YJW.

References

1. Rysanek D, Zouharova M, Babak V. Monitoring major mastitis pathogens at the population level based on examination of bulk tank milk samples. *J Dairy Res.* 2009; 76(1):117–23. doi: PMID: [19121238](#)
2. Makovec JA, Ruegg PL. Results of milk samples submitted for microbiological examination in Wisconsin from 1994 to 2001. *J Dairy Sci.* 2003; 86(11):3466–72. PMID: [14672176](#)
3. Katholm J, Bennedsgaard TW, Koskinen MT, Rattenborg E. Quality of bulk tank milk samples from Danish dairy herds based on real-time polymerase chain reaction identification of mastitis pathogens. *J Dairy Sci.* 2012; 95(10):5702–8. doi: PMID: [22921631](#)
4. Bradley AJ. Bovine mastitis: An evolving disease. *Vet J.* 2002; 164(2):116–28. PMID: [12359466](#)
5. Jayarao BM, Pillai SR, Sawant AA, Wolfgang DR, Hegde NV. Guidelines for monitoring bulk tank milk somatic cell and bacterial counts. *J Dairy Sci.* 2004; 87(10):3561–73. PMID: [15377636](#)
6. Phuektes P, Browning CF, Anderson G, Mansell PD. Multiplex polymerase chain reaction as a mastitis screening test for *Staphylococcus aureus*, *Streptococcus agalactiae*, *Streptococcus dysgalactiae* and *Streptococcus uberis* in bulk milk samples. *J Dairy Res.* 2003; 70(2):149–55. PMID: [12800868](#)
7. Riekerink R, Barkema HW, Scholl DT, Poole DE, Kelton DF. Management practices associated with the bulk-milk prevalence of *Staphylococcus aureus* in Canadian dairy farms. *Prev Vet Med.* 2010; 97(1):20–8. doi: PMID: [20696486](#)
8. Godkin MA, Leslie KE. Culture of bulk tank milk as a mastitis screening test: A brief review. *Can Vet J.* 1993; 34(10):601–5. PMID: [17424304](#)
9. Bramley AJ, Mckinnon CH, Staker RT, Simpkin DL. The effect of udder infection on the bacterial-flora of the bulk milk of 10 dairy herds. *J Appl Bacteriol.* 1984; 57(2):317–23. PMID: [6389466](#)
10. Riekerink RGO, Barkema HW, Veenstra S, Poole DE, Dingwell RT, Keefe GP. Prevalence of contagious mastitis pathogens in bulk tank milk in Prince Edward Island. *Can Vet J.* 2006; 47(6):567. PMID: [16808229](#)
11. Jayarao BM, Wolfgang DR. Bulk-tank milk analysis: A useful tool for improving milk quality and herd udder health. *Vet Clin N Am-Food A.* 2003; 19(1):75–92.
12. Syring C, Boss R, Reist M, Bodmer M, Hummerjohann J, Gehrig P, et al. Bovine mastitis: The diagnostic properties of a PCR-based assay to monitor the *Staphylococcus aureus* genotype B status of a herd, using bulk tank milk. *J Dairy Sci.* 2012; 95(7):3674–82. doi: PMID: [22720925](#)
13. Viguier C, Arora S, Gilmartin N, Welbeck K, O Kennedy R. Mastitis detection: current trends and future perspectives. *Trends Biotechnol.* 2009; 27(8):486–93. doi: PMID: [19616330](#)
14. Koskinen MT, Holopainen J, Pyörälä S, Bredbacka P, Pitkälä A, Barkema HW, et al. Analytical specificity and sensitivity of a real-time polymerase chain reaction assay for identification of bovine mastitis pathogens. *J Dairy Sci.* 2009; 92(3):952–9. doi: PMID: [19233788](#)
15. Mweu MM, Toft N, Katholm J, Nielsen SS. Evaluation of two herd-level diagnostic tests for *Streptococcus agalactiae* using a latent class approach. *Vet Microbiol.* 2012; 159(1–2):181–6. doi: PMID: [22542270](#)
16. Pulido-Landinez M, Laviniki V, Sanchez-Ingunza R, Guard J, Do Nascimento VP. Use of FTA cards for the transport of DNA samples of *Salmonella* spp. from poultry products from Southern Brazil. *Acta Sci Vet.* 2012; 40(10734).
17. Rajendram D, Ayenza R, Holder FM, Moran B, Long T, Shah HN. Long-term storage and safe retrieval of DNA from microorganisms for molecular analysis using FTA matrix cards. *J Microbiol Meth.* 2006; 67(3):582–92.
18. Whatman. FTA Nucleic Acid Collection, Storage and Purification. 2009. Available:

19. Linhares DC, Rovira A, Torremorell M. Evaluation of Flinders Technology Associates cards for collection and transport of samples for detection of Porcine reproductive and respiratory syndrome virus by reverse transcription polymerase chain reaction. *J Vet Diagn Invest.* 2012; 24(2):328–32. doi: PMID: [22362527](#)
20. Abdelwhab EM, Lueschow D, Harder TC, Hafez HM. The use of FTA (R) filter papers for diagnosis of avian influenza virus. *J Virol Methods.* 2011; 174(1–2):120–2. doi: PMID: [21419801](#)
21. Ahmed HA, MacLeod ET, Hide G, Welburn SC, Picozzi K. The best practice for preparation of samples from FTA (R) cards for diagnosis of blood borne infections using African trypanosomes as a model system. *Parasite Vector.* 2011; 4(68).
22. Grund E, Darissa O, Adam G. Application of FTA (R) cards to sample microbial plant pathogens for PCR and RT-PCR. *J Phytopathol.* 2010; 158(11–12):750–7.
23. Stangegaard M, Borsting C, Ferrero-Miliani L, Frank-Hansen R, Poulsen L, Hansen AJ, et al. Evaluation of four automated protocols for extraction of DNA from FTA cards. *JALA-J Lab Autom.* 2013; 18(5):404–10.
24. Tilsala-Timisjärvi A, Alatossava T. Rapid DNA preparation from milk and dairy process samples for the detection of bacteria by PCR. *Food Microbiol.* 2004; 21(3):365–8.
25. Kraus RHS, van Hooff P, Waldenstrom J, Latorre-Margalef N, Ydenberg RC, Prins HHT. Avian influenza surveillance: on the usability of FTA (R) cards to solve biosafety and transport issues. *Wildfowl.* 2009(Sp. Iss. 2.):215–23.
26. Taponen S, Salmikivi L, Simojoki H, Koskinen MT, Pyörala S. Real-time polymerase chain reaction-based identification of bacteria in milk samples from bovine clinical mastitis with no growth in conventional culturing. *J Dairy Sci.* 2009; 92(6):2610–7. doi: PMID: [19447993](#)
27. Mahmmod YS, Toft N, Katholm J, Gronbaek C, Klaas IC. Estimation of test characteristics of real-time PCR and bacterial culture for diagnosis of subclinical intramammary infections with *Streptococcus agalactiae* in Danish dairy cattle in 2012 using latent class analysis. *Prev Vet Med.* 2013; 109(3–4):264–70. doi: PMID: [23194895](#)
28. Koskinen MT, Wellenberg GJ, Sampimon OC, Holopainen J, Rothkamp A, Salmikivi L, et al. Field comparison of real-time polymerase chain reaction and bacterial culture for identification of bovine mastitis bacteria. *J Dairy Sci.* 2010; 93(12):5707–15. doi: PMID: [21094742](#)
29. Nielsen PK, Petersen MB, Nielsen LR, Halasa T, Toft N. Latent class analysis of bulk tank milk PCR and ELISA testing for herd level diagnosis of *Mycoplasma bovis*. *Prev Vet Med.* 2015; 121(3–4):338–42. doi: PMID: [26342789](#)
30. Cicconi-Hogan KM, Gamroth M, Richert R, Ruegg PL, Stiglbauer KE, Schukken YH. Associations of risk factors with somatic cell count in bulk tank milk on organic and conventional dairy farms in the United States. *J Dairy Sci.* 2013; 96(6):3689–702. doi: PMID: [23548286](#)
31. Fenlon DR, Logue DN, Gunn J, Wilson J. A study of mastitis bacteria and herd management practices to identify their relationship to high somatic cell counts in bulk tank milk. *Br Vet J.* 1995; 151(1):17–25. PMID: [7735867](#)
32. Barkema HW, Schukken YH, Lam T, Beiboer ML, Wilmink H, Benedictus G, et al. Incidence of clinical mastitis in dairy herds grouped in three categories by bulk milk somatic cell counts. *J Dairy Sci.* 1998; 81(2):411–9. PMID: [9532494](#)
33. Wilson DJ, Gonzale RN, Sears PM. Segregation or use of separate milking units for cows infected with *Staphylococcus aureus*: Effects on prevalence of infection and bulk tank somatic cell count. *J Dairy Sci.* 1995; 78(9):2083–5. PMID: [8550917](#)
34. Koop G, Dik N, Nielen M, Lipman L. Short communication: Repeatability of differential goat bulk milk culture and associations with somatic cell count, total bacterial count, and standard plate count. *J Dairy Sci.* 2010; 93(6):2569–73. doi: PMID: [20494165](#)
35. Klossowska A, Malinowski E, Kuzma K. Relationship between somatic cell counts in cow quarter fore-milk samples and etiological agents of mastitis. *Med Weter.* 2005; 61(1):53–7.
36. Malinowski E, Lassa H, Klossowska A, Markiewicz H, Kaczmarowski M, Smulski S. Relationship between mastitis agents and somatic cell count in foremilk samples. *B Vet I Pulawy.* 2006; 50(3):349–52.
37. Berchtold B, Bodmer M, van den Borne B, Reist M, Graber HU, Steiner A, et al. Genotype-specific risk factors for *Staphylococcus aureus* in Swiss dairy herds with an elevated yield-corrected herd somatic cell count. *J Dairy Sci.* 2014; 97(8):4886–96. doi: PMID: [24881801](#)
38. Dinsmore RP. Biosecurity for mammary diseases in dairy cattle. *Vet Clin N Am-Food A.* 2002; 18(1):115.
39. Riekerink R, Barkema HW, Kelton DF, Scholl DT. Incidence rate of clinical mastitis on Canadian dairy farms. *J Dairy Sci.* 2008; 91(4):1366–77. doi: PMID: [18349229](#)

40. Zadoks RN, Gillespie BE, Barkema HW, Sampimon OC, Oliver SP, Schukken YH. Clinical, epidemiological and molecular characteristics of *Streptococcus uberis* infections in dairy herds. *Epidemiol Infect.* 2003; 130(2):335–49. PMID: [12729202](#)
41. Brito M, Brito J, de Souza HM, Vargas OL. Evaluation of the sensitivity of bulk tank milk cultures for the isolation of contagious bovine mastitis pathogens. *Pesqui Vet Brasil.* 1998; 18(1):39–44.